Proteins

Screening Libraries

Product Data Sheet

MR 409

Molecular Weight:

Cat. No.: HY-P3304 CAS No.: 1445155-39-4 Molecular Formula: $C_{153}H_{252}N_{44}O_{43}$

3395.91

Sequence Shortening: {N-Me}-Y-{D-Ala}-DAIFTNSYR-{Orn}-VL-{Abu}-QLSAR-{Orn}-LLQDI-{Nle}-DR-NHMe

Target: Others Others Pathway:

Storage: Sealed storage, away from moisture and light, under nitrogen

> 2 years Powder -80°C -20°C 1 year

* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture

and light, under nitrogen)

BIOLOGICAL ACTIVITY

Description

MR 409 is a selected growth hormone-releasing hormone (GHRH) agonist. MR 409 has remarkable neuroprotective effects through enhancing endogenous neurogenesis in cerebral ischemic mice. MR 409 also inhibits the in vivo growth of lung cancer^{[1][2][3][4]}.

In Vitro

One week post-myocardial infarction (MI), MR 409 (MR-409) significantly reduces plasma levels of IL-2, IL-6, IL-10 and TNF- α compared to placebo^[1].

MR 409 (MR-409; 1 μM) suppresses p53 expression in bovine pulmonary arterial endothelial cells (BPAECs)^[2].

MR 409 (1 $\mu\text{M})$ induces the activation of JAK2, STAT3 and ERK1/2 $^{[2]}.$

MR 409 (MR-409; 1 and 5 µM) decreases LPS-induced PGE₂ and 8-iso-PGF_{2a} levels, in a dose-dependent manner^[3].

MR 409 (1 and 5µM) decreases LPS-induced lactate dehydrogenase (LDH) activity and nitrite production, without showing a dose-dependent effect^[3].

MR 409 (1 and 5 µM) decreases LPS-induced gene expression of COX-2, NF-κB and iNOS in colon specimens, without a dosedependent effect^[3].

MR 409 (MR-409) can stimulate endogenous neurogenesis and improve the tMCAO-induced loss of neuroplasticity. MR 409 also enhances the proliferation and inhibits apoptosis of neural stem cells treated with oxygen and glucose deprivation-

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Western Blot Analysis^[2]

Cell Line:	BPAEC
Concentration:	1 μΜ
Incubation Time:	24, 48 and 72 hours
Result:	Significantly suppressed p53 expression levels after 48 and 72 hours of treatment.

In Vivo

MR 409 (MR-409) inhibits the in vivo growth of lung cancer xenografted into nude mice [3].

MR 409 (5 μ g; s.c. treated daily for 4 weeks) reduces nociceptive response in mice^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	Adult C57/BL6 male mice (5 weeks old, weight 20-22 g) ^[3]
Dosage:	5 μg
Administration:	Treated daily by subcutaneous (s.c.) administration of 0.1 mL solution
Result:	Showed maximal antinociceptive effects at 2 weeks. Sustained a moderate analgesic effect at 4 weeks.

REFERENCES

- [1]. Rosemeire M Kanashiro-Takeuchi, et al. New therapeutic approach to heart failure due to myocardial infarction based on targeting growth hormone-releasing hormone receptor. Oncotarget. 2015;6(12):9728-39.
- [2]. Mohammad A Uddin, et al. GHRH antagonists support lung endothelial barrier function. Tissue Barriers. 2019;7(4):1669989.
- [3]. Lucia Recinella, et al. Protective effects of growth hormone-releasing hormone analogs in DSS-induced colitis in mice. Sci Rep. 2021 Jan 28;11(1):2530.
- [4]. Yueyang Liu, et al. Agonistic analog of growth hormone-releasing hormone promotes neurofunctional recovery and neural regeneration in ischemic stroke. Proc Natl Acad Sci U S A. 2021 Nov 23;118(47):e2109600118.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$

Tel: 609-228-6898

Fax: 609-228-5909

 $\hbox{E-mail: } tech @ {\tt MedChemExpress.com}$

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA