

# **Product** Data Sheet

## **DBCO-amine**

Cat. No.:HY-W000423CAS No.:1255942-06-3Molecular Formula: $C_{18}H_{16}N_2O$ Molecular Weight:276.33Target:ADC Linker

Pathway: Antibody-drug Conjugate/ADC Related

**Storage:** -20°C, protect from light

\* In solvent: -80°C, 6 months; -20°C, 1 month (protect from light)

### **SOLVENT & SOLUBILITY**

In Vitro

DMSO: 100 mg/mL (361.89 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	3.6189 mL	18.0943 mL	36.1886 mL
	5 mM	0.7238 mL	3.6189 mL	7.2377 mL
	10 mM	0.3619 mL	1.8094 mL	3.6189 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (9.05 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (9.05 mM); Clear solution

### **BIOLOGICAL ACTIVITY**

Description	DBCO-amine is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs) <sup>[1]</sup> .				
IC <sub>50</sub> & Target	Cleavable				
In Vitro	ADCs are comprised of an antibody to which is attached an ADC cytotoxin through an ADC linker.  MCE has not independently confirmed the accuracy of these methods. They are for reference only.				

#### **REFERENCES**

1]. Ermira Pazolli, et al. Splicing modulator antibody-drug conjugates and methods of use. WO2019232449A1.							
	Caution: Product has not l	been fully validated for mo	edical applications. For researd	-h use only			
	Tel: 609-228-6898	Fax: 609-228-5909	E-mail: tech@MedChemE				
	Address: 1 De	er Park Dr, Suite Q, Monmo	outh Junction, NJ 08852, USA				

Page 2 of 2 www.MedChemExpress.com