


Lurbinectedin

Cat. No.:	HY-16293
CAS No.:	497871-47-3
Molecular Formula:	C ₄₁ H ₄₄ N ₄ O ₁₀ S
Molecular Weight:	784.87
Target:	DNA/RNA Synthesis; DNA Alkylator/Crosslinker
Pathway:	Cell Cycle/DNA Damage
Storage:	Powder -20°C 3 years In solvent -80°C 6 months -20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

DMSO : 20 mg/mL (25.48 mM; Need ultrasonic)

	Solvent Concentration	Mass	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM		1.2741 mL	6.3705 mL	12.7410 mL
	5 mM		0.2548 mL	1.2741 mL	2.5482 mL
	10 mM		0.1274 mL	0.6370 mL	1.2741 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Lurbinectedin (PM01183) is a DNA minor groove covalent binder with potent anti-tumour activity; inhibits RMG1 and RMG2 cell growth with IC₅₀ values of 1.25 and 1.16 nM, respectively.

IC₅₀ & Target

IC₅₀: 1.25 nM (RMG1), 1.16 nM (RMG2)^[1]

In Vitro

PM01183 is a new synthetic tetrahydroisoquinoline alkaloid that is currently in phase I clinical development for the treatment of solid tumours. PM01183–DNA adducts in living cells give rise to double-strand breaks, triggering S-phase accumulation and apoptosis. The potent cytotoxic activity of PM01183 is ascertained in a 23-cell line panel with a mean GI₅₀ value of 2.7 nM^[2]. Lurbinectedin exhibits significant antitumor activity toward chemosensitive and chemoresistant human ovarian clear cell carcinoma (CCC) cells in vitro^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Mouse CCC cell xenografts reveals that lurbinectedin significantly inhibits tumor growth. Lurbinectedin plus SN-38 results in a significant synergistic effect^[1]. In four murine xenograft models of human cancer, PM01183 inhibits tumour growth significantly with no weight loss of treated animals^[2]. Single lurbinectedin or NSC 119875-combined therapies are effective in treating NSC 119875-sensitive and NSC 119875-resistant preclinical ovarian tumor models. The strongest synergistic

effect is observed for combined treatments, especially in NSC 119875-resistant tumors. Lurbinectedin tumor growth inhibition is associated with reduced proliferation, increased rate of aberrant mitosis, and subsequent induced apoptosis^[3]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay ^[1]

The MTS assay is used to analyze the effects of each drug. Cells are plated in 96-well plates and treated with PM01183 (0, 0.1, 0.3, 1, 3 nM) . After 48 hours' incubation, the number of surviving cells is assessed by determining the A490nm of the dissolved formazan product after the addition of MTS for 1 hour. Cell viability is calculated as follows: Aexp group / Acontrol×100^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[3]

Mice: Mice are transplanted with fragments of OVAX1 and OVAX1R tumors, and when tumors reaches a homogeneous palpable size are randomly allocated into the treatment groups: i) Placebo; ii) Lurbinectedin (0.18 mg/kg); iii) NSC 119875 (3.5 mg/kg); and iv) Lurbinectedin plus NSC 119875 (0.18 + 3.5 mg/kg). Drugs are i.v. administered once per week for 3 consecutive weeks (days 0, 7, and 14). Seven days after the final dose (day 21), animals are sacrificed, their ovaries dissected out, and weighed. Representative fragments are either frozen in nitrogen or fixed and then processed for paraffin embedding^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Thorac Cancer. 2022 Apr;13(8):1184-1192.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Takahashi R, et al. Preclinical Investigations of PM01183 (Lurbinectedin) as a Single Agent or in Combination with Other Anticancer Agents for Clear Cell Carcinoma of the Ovary. PLoS One. 2016 Mar 17;11(3):e0151050.
- [2]. Leal JF, et al. PM01183, a new DNA minor groove covalent binder with potent in vitro and in vivo anti-tumour activity. Br J Pharmacol. 2010 Nov;161(5):1099-110.
- [3]. Vidal A, et al. Lurbinectedin (PM01183), a new DNA minor groove binder, inhibits growth of orthotopic primary graft of NSC 119875-resistant epithelial ovarian cancer. Clin Cancer Res. 2012 Oct 1;18(19):5399-411.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA